

## MEDICATION ORDERS-KRYSTEXXA (PEGLOTICASE)

PATIENT INFORMATION	
Name:	DOB:
Allergies:	Date of Referral:

REFERRAL STATUS
<input type="checkbox"/> New Referral <input type="checkbox"/> Dose or Frequency Change <input type="checkbox"/> Order Renewal

INFUSION OFFICE PREFERENCES (Optional)
Preferred Location*:

\*List of infusion center locations may be found at: <https://metroinfusioncenter.com/infusion-center-locations/>  
 Please note: Requests will be accommodated based on infusion center availability and are not guaranteed.

DIAGNOSIS AND ICD 10 CODE	
<input type="checkbox"/> Chronic gout with Tophus	ICD 10 Code: M1A.9xx1
<input type="checkbox"/> Chronic gout without Tophus	ICD 10 Code: M1A.9XX0

REQUIRED DOCUMENTATION	
<input type="checkbox"/> This signed order form by the provider <input type="checkbox"/> Patient demographics AND insurance information <input type="checkbox"/> Uric acid level	<input type="checkbox"/> Clinical/Progress notes <input type="checkbox"/> Labs and Tests supporting primary diagnosis <input type="checkbox"/> G6PD test results
List Tried & Failed Therapies:	
1)	
2)	
3)	

MEDICATION ORDERS	
Dosing	<input type="checkbox"/> Krystexxa 8mg IV every 2 weeks
Refills:	<input type="checkbox"/> X 6 months <input type="checkbox"/> X 1 year <input type="checkbox"/> ____ doses

PREMEDICATIONS
<input type="checkbox"/> Acetaminophen 650mg PO prior to Krystexxa infusion <input type="checkbox"/> Diphenhydramine 25mg PO prior to Krystexxa infusion <input type="checkbox"/> Methylprednisolone 40mg Slow IV Push prior to Krystexxa infusion <input type="checkbox"/> Other:

Please note: if an infusion reaction occurs, the on-call physician will order appropriate rescue medications as deemed medically necessary. This may also include pausing, reducing the rate of infusion or discontinuing the medication.

PRESCRIBER INFORMATION		
Prescriber Name:		
Office Phone:	Office Fax:	Office Email:
Prescriber Signature:		Date: