Pemgarda (pemivibart)



Infusion Office Preference:			
PATIENT INFORMATION			
Date: Patient Name:		DOB:	
□ NKDA Allergies:			
PROVIDER INFORMATION			
Office Contact Name:	Office Email:		
Prescribing Providers Name:	Provider NPI:		
Office Address:	City:	State:	Zip:
Office Phone Number:	Office Fax Number:		
DIAGNOSIS /ICD 10			
medications or treatments and are unlikely to mount an adequate immune response to COVID-19 vaccination and : Not currently infected with SARS-CoV-2 and who have not had a known recent exposure to an individual with SARS-CoV-2 (ICD 10 for this diagnosis is Q0224)			
REQUIRED DOCUMENTATION/Testing			
 □ This signed order form by the provider □ Patient demographics AND insurance information 	☐ Clinical/Progress notes supporting diagnosis above☐ Verification that patient has not had severe		
	-	hypersensitivity reaction to a COVID-19 vaccine	
MEDICATION ORDERS			
□ 4500mg iv one-time infusion over 60 minutes using 0.2micron inline filter			
□ 4500mg iv every 3 months over 60 minutes using 0.2micron inline filter			
Patients must be monitored for 2 hours after the infusion.			
Provider Name (Print) Physician Sign	ature:	n	late: