

Leqembi (lecanemab-irmb)

Infusion Office Preference: _____

PATIENT INFORMATION		
Date:	Patient Name:	DOB:
☐ NKDA Allergies:	Weight (lbs / kg):	Height:
Patient Status: ☐ New to Therapy ☐ Continuing Therapy - Last Treatment Date:		Next Due Date:
DIAGNOSIS AND ICD 10 CODE		
☐ Alzheimer's disease with early onset	ICD 10 Code: G30.0	
☐ Mild Cognitive Impairment, So stated	ICD 10 Code: G31.84	
☐ other ICD10 Code:)	Description:	
G30.X CODES BELOW REQUIRE SECONDARY F02.8x DIAGNOSIS CODE- PLEASE SELECT ONE FROM EACH COLUMN		
		<u>Secondary</u>
☐ G30.1 Alzheimer's disease late onset	☐ F02.80 Dementia without behavioral disturbance	
☐ G30.8 Other Alzheimer's disease	☐ F02.81 Dementia with behavioral disturbance	
☐ G30.9 Alzheimer's disease , unspecified		
REQUIRED DOCUMENTATION		
☐ This signed order form by the provider ☐ Patient demographics AND insurance info ☐ Clinical/Progress notes		
Prescriber must indicate that the following requirements have been met (provide supporting documentation)		
☐ Beta Amyloid Pathology Confirmed via: ↳ ☐ Amyloid PET Scan OR ☐ CFS Analysis-Date: _____ Result: _____		
☐ Cognitive Assessment Used: _____ Date: _____ Result: _____		
☐ ApoE εε4 Genetic Test - Date: _____ Result: ☐ Homozygote ☐ Heterozygote ☐ Noncarrier		
MEDICARE REGISTRATION # IF APPLICABLE _____		
MEDICATION ORDERS(Note: Only one stage of treatment may be ordered at a time)		
☐ Stage 1 (Infusions #1 and #2) ✓Leqembi 10mg/kg IV every two weeks x 2 doses. Each infusion to be given over one hour. Required Documentation to Initiate this Phase: MRI of brain within one year prior to first infusion. Date of MRI: _____		
☐ By checking this box, I confirm that Beta Amyloid Pathology has been confirmed via CSF or PET.		
☐ Stage 2 (Infusions #3 and #4) ✓Leqembi 10mg/kg IV every two weeks x 2 doses. Each infusion to be given over one hour. Required Documentation to Initiate this Phase: ☐ By checking this box, I confirm that patient has undergone MRI of brain before dose #3. I have reviewed the results and clear patient to proceed with infusions #3 and #4		
☐ Stage 3 (Infusions #5 and #6) ✓Leqembi 10mg/kg IV every two weeks x 2 doses. Each infusion to be given over one hour. Required Documentation to Initiate this Phase: ☐ By checking this box, I confirm that patient has undergone MRI of brain before dose #5. I have reviewed the results and clear patient to proceed with infusions #5 and #6.		
☐ Stage 4 (Infusions #7-13) ✓Leqembi 10mg/kg IV every two weeks x 7 doses. Each infusion to be given over one hour. Required Documentation to Initiate this Phase: ☐ By checking this box, I confirm that patient has undergone MRI of brain before dose #7. I have reviewed the results and clear patient to proceed with infusions #7 through #13.		
* Stage 5 (Infusions #14 & beyond) MUST SELECT ONE DOSING BELOW ☐ Leqembi 10mg/kg IV every two weeks over one hour OR ☐ Leqembi 10mg/kg IV every four weeks over one hour. Required Documentation to Initiate this Phase: ☐ By checking this box, I confirm that patient has undergone MRI of brain before dose #14. I have reviewed the results and clear patient to proceed with infusions #14 and beyond as ordered above		

Fax referral to 866-507-1164 or email to bionurses@metroinfusioncenter.com

All information contained in this order form is strictly confidential and will become 01/29/25part of the patient's medical record.

Revised 9/15/25

PRE-INFUSION :

- ☒ Confirm baseline MRI results prior to initiation of treatment
- ☒ Confirm MRI completed and reviewed by prescriber prior to the 5th, 7th, and 14th treatment
- ☒ Measure and record weight prior to each treatment to determine dose
- ☒ Hold infusion and notify provider if patient reports:
 - Headache
 - Dizziness
 - Nausea
 - Vision Changes
 - New or worsening confusion

Post -INFUSION :

- ☒ Educate patient/care partner to report headache, dizziness, nausea, vision changes, or new/worsening confusion.
- ☒ Fax infusion record to provider below :

PRESCRIBER INFORMATION

Provider Name (print)

Provider Signature**:

Date:

Office Phone

Office Fax

***Amyloid Related Imaging Abnormalities (ARIA): Enhanced clinical vigilance for ARIA is recommended during the first 14 weeks of treatment with LEQEMBI. Referring physician to monitor patient*

Fax referral to 866-507-1164 or email to bionurses@metroinfusioncenter.com

All information contained in this order form is strictly confidential and will become 01/29/25 part of the patient's medical record.

Revised 9/15/25