



# METRO INFUSION CENTER

## Octreotide Acetate (Sandostatin LAR)

Name: \_\_\_\_\_

DOB: \_\_\_\_\_

Diagnosis/Code: \_\_\_\_\_ / \_\_\_\_\_

Cancer Stage: \_\_\_\_\_

**Dose calculation:**

Flat dose, not a weight based medication

**Dosing Guidelines/Parameters:** *Provider must select hold parameters that will trigger a call from the Infusion Staff***Hydration Orders:**

Not required

**Premedication and Antiemetic Orders:**

Not required

**Treatment Orders:**

DRUG	DOSE	ROUTE	DAYS TO BE GIVEN
<input type="checkbox"/> Octreotide Acetate (Sandostatin LAR)	10mg	IM*	Every 28 days
<input type="checkbox"/> Octreotide Acetate (Sandostatin LAR)	20mg	IM*	Every 28 days
<input type="checkbox"/> Octreotide Acetate (Sandostatin LAR)	30mg	IM*	Every 28 days
<input type="checkbox"/> Octreotide Acetate (Sandostatin LAR)	40mg	IM*	Every 28 days

\* Give in the outer gluteal region with recommended needle size for administration of SANDOSTATIN LAR DEPOT is the 1½" 19-gauge safety injection needle (supplied in the drug product kit). For patients with a greater skin to muscle depth, a size 2" 19-gauge needle (not supplied) may be used

Date of intended first treatment at Metro Infusion Center: \_\_\_\_\_

Subsequent treatment may be given +/- 2 days or as otherwise specified:

***This order is good for 1 year from the date ordered***

Other:

Call referring provider for:

Date:

Referring Provider: \_\_\_\_\_ Phone# \_\_\_\_\_

SIGNATURE REQUIRED

PRINTED NAME REQUIRED

**Office Contact name/number:**

All information in this order is strictly confidential and will become part of the patient's medical record. Contact us with questions at (877)448-3627. Send completed form and all documentation to confidential email:

[MICreferral@metroinfusioncenter.com](mailto:MICreferral@metroinfusioncenter.com) or fax to (866)507-1164

Revised 12/2/25